arXiv:1511.07352v1 [astro-ph.SR] 23 Nov 2015

Probing The Accreting Hot Components in Six S-Type Symbiotic Variables

Edward M. Sion

Department of Astrophysics & Planetary Science, Villanova University, 800 Lancaster Avenue, Villanova, PA 19085, USA

edward.sion@villanova.edu

Patrick Godon

Department of Astrophysics & Planetary Science, Villanova University, 800 Lancaster Avenue, Villanova, PA 19085, USA

patrick.godon@villanova.edu

Joanna Mikolajewska

Copernicus Astronomical Center, Warsaw, Poland

mikolaj@camk.edu.pl

Bassem Sabra

Dept. of Physics & Astronomy, Notre Dame University, Louaize; Zouk Mosbeh, Lebanon

bsabra@ndu.edu.lb

Craig Kolobow

Department of Physics & Astronomy, Florida Institute of Technology, Melbourne, FL 98195, USA

craig.kolobow@villanova.edu

ABSTRACT

We have carried out a spectroscopic analysis of the far ultraviolet spectra of six symbiotic variables. Two systems, LT Del, which has had one recorded outburst, and BD-21 3873 (= IV Vir) which has had no recorded outburst, are

yellow symbiotic systems. Two other systems, V443 Her and RW Hya, have also never had a recorded outburst. Two other symbiotics, StHa190 and CQ Dra, are more strongly interacting with an outburst history. We have studied these systems during their quiescence in order to shed light on the nature of their hot components by fitting their archival far ultraviolet spectra with optically thick accretion disk models and NLTE model white dwarf photospheres. Using the critical advantage offered by extending wavelength coverage down to the Lyman Limit with FUSE spectra, we find that the hot component in RW Hya is a low mass white dwarf with a surface temperature of 160,000K while the symbiotic system CQ Dra is a triple system with a red giant transferring matter to a hot component made up of a cataclysmic variable whose white dwarf has a surface temperature of ∼50,000K. Implications are discussed.

Subject headings: stars: binaries: stars: symbiotics: individuals: IV Vir, StHa190, CQ Dra, RW Hya, LT Del, V443 Her

1. Introduction

Up to now, virtually all of the temperatures and luminosities of symbiotic hot components have been derived using the modified Zanstra method to obtain radiation temperatures needed to account for photoionization of the observed emission lines (Muerset et al. 1991; Muerset & Nussbaumer 1994; Skopal 2005), or by simply assuming a photospheric temperature. This is understandable given the daunting complexity of symbiotic stars and the possible sources of FUV continuum radiation that may arise from that complexity. Nevertheless, many models have been proposed for the origin of the emission lines including the assumption that they all arise from photoionization. Moreover, the temperatures are derived for the most part from the He II (1640) recombination line and the H lines, but they yield different temperatures with the temperature from the He II lines much higher than that from the H lines. He II lines arise from a wind or from wind collision shocks or coronae? Added to this uncertainty is the fact that for the very few systems analyzed up to now with actual NLTE photospheric models (mostly post-novae), the model-derived temperatures for hot accreting white dwarfs are a factor of two to three lower than the temperatures derived by Zanstra techniques.

By far the most extensive FUV database of symbiotic systems is The International Ultraviolet Explorer (IUE) spacecraft. Among the approximately two dozen systems in the IUE archive with steeply rising FUV continua, we selected six systems for this study; V443 Her, LT Del, BD-21 3873, StHa190, CQ Dra and RW Hya. Two of the systems, RW Hya and

CQ Dra, have usable archival FUSE spectra as well, while RW Hya was observed with the Hubble Space Telescope and has HST spectra in the MAST archive. Two systems, LT Del, which has had one recorded outburst, and BD-21 3873 (= IV Vir) which has had no recorded outburst, are both yellow symbiotic systems. They are metal-poor, and their red giants are s-process enhanced. This s-process enhancement is due to past mass transfer (when the present WD was on AGB), and provides very strong indications that the WD mass should be at least $\sim 0.55 M_{\odot}$. Two systems, V443 Her and RW Hya, have also never had a recorded outburst. Two other symbiotics, StHa190 and CQ Dra, are more strongly interacting with an outburst history.

Three of these six systems show ellipsoidal light curves, especially in the red and near-IR range (where the red giant dominates the spectrum), which strongly indicates that the red giant is filling or nearly filling its ROCHE lobe. Therefore the mass transfer and accretion is mostly though L1, and hence an accretion disk is highly likely to be present. Thus, Their mass transfer/accretion rates should also be relatively high, $\sim 10^{-7} M_{\odot}$/yr, as the numerical hydrodynamic simulations results on mass transfer in such systems

The published orbital and physical parameters of these six systems are given in Table 1 where we list, by row: (1) the system name; (2) orbital period in days P_{orb}; (3) the distance estimate in pc; (4) the orbital inclination in degrees; (5) estimated mass of the hot component, M_{hot}; (6) the orbital separation in A.U. and (7) the color excess E(B-V).

We studied these systems during their quiescence in order to shed light on the nature of their hot components by fitting the observations with optically thick accretion disk models and model white dwarf photospheres. Our aim is to determine for the first time, with NLTE model accretion disks and model photospheres, the accretion rates, accretion efficiency and the temperatures and luminosities of the hot components and ascertain whether their FUV radiation is from accretion light or WD photospheric light or both.

In the sections below, we present: (1) an observing log of the archival FUV spectra utilized in our study; (2) describe our model accretion disks, high gravity photospheres and modeling analysis techniques; (3) describe each system in greater depth and summarize our model fitting results. In section 4, we state our conclusions.

2. Archival Far Ultraviolet Spectroscopic Observations

Each far UV spectrum was downloaded from the NEWSIPS International Ultraviolet Explorer (IUE) archive, the Far Ultraviolet Spectroscopic Explorer archive and, for RW Hya, the Hubble Space Telescope archive on MAST. Only one usable IUE spectrum was available

for LT Del, V443 Her, and StHa190, two IUE spectra were available for BD-21 3873 and CQ Dra while several IUE spectra were available for RW Hya. All spectra were taken during quiescence.

Table 2 provides the IUE observing log for all six systems where we list the observing details for each system, by row: (1) the system name; (2) the SWP spectrum No.; (3) the date of the IUE observation; (4) the exposure time in seconds; (5) the continuum counts and; (6) the background counts.

Archival FUSE spectra were available for CQ Dra and RW Hya. For CQ Dra, FUSE spectrum #D9050401000, with 10 exposures, was obtained on January 9, 2004 through the LWRS aperture with an exposure time of 36719s, and a central wavelength of 1046A. For RW Hya, FUSE spectrum #E1460103000, with 4 exposures, was obtained on July 21, 2004 through the LWRS aperture with an exposure time of 4353s, and a central wavelength of 1046A.

An HST spectrum of RW Hydrae was obtained with the Goddard High Resolution Spectrograph (GHRS) in 1996 July 21, in ACCUM mode with the 2.0 x 2.0 aperture. The G140L disperser was used, giving a wavelength coverage of 1150A to 1700A. The spectrum was obtained at approximately orbital quadrature, phase 0.72 with an exposure time of 872s.

3. Model Accretion Disks and model Photospheres

We adopted model accretion disks from the optically thick disk model grid of Wade & Hubeny (1998). In these accretion disk models, the outermost disk radius, R_{out}, is chosen so that $T_{eff}(R_{out})$ is near 10,000K since disk annuli beyond this point, which are cooler zones with larger radii, would provide only a very small contribution to the mid and far UV disk flux, particularly the SWP FUV band pass. The mass transfer rate is assumed to be the same for all radii.

Thus, the run of disk temperature with radius is taken to be:

$$T_{eff}(r) = T_s x^{-3/4}(1 - x^{-1/2})^{1/4} \tag{1}$$

where $x = r/R_{wd}$ and $\sigma T_s^4 = 3GM_{wd}\dot{M}/8\pi R_{wd}^3$

Limb darkening of the disk is fully taken into account in the manner described by Diaz et al. (1996) involving the Eddington-Barbier relation, the increase of kinetic temperature with depth in the disk, and the wavelength and temperature dependence of the Planck

function.

For a given spectrum, we carry out fits for every combination of $\dot{M}$, inclination and white dwarf mass in the Wade & Hubeny (1998) library.

The values of i are 18, 41, 60, 75 and 81. The range of accretion rates covers $-10.5 < log(\dot{M}) < -8.0$ in steps of 0.5 in the log and five different values of the white dwarf mass, namely, 0.35, 0.55, 0.8, 1.03, and 1.21 $M_{\odot}$.

For the disk models, we ran every combination of inclination angle i = 18, 41, 60, 75, 81 degrees, M_{wd}= 0.35, 0.55, 0.80, 1.03, and 1.21 $M_{\odot}$ and $Log(\dot{M})$= -8.0, -8.5, -9.0, -9.5, -10.0, -10.5. Theoretical, high gravity, solar composition photospheric spectra were computed by first using the code TLUSTY version 200 Hubeny (1988) to calculate the atmospheric structure and then SYNSPEC version 48 (Hybeny & Lanz 1995) to construct synthetic spectra. We compiled a library of photospheric spectra covering the temperature range from 13,000K to 80,000K in increments of 500K (for low T_{wd}), 1000K (for intermediate T_{wd}) and 5000K (for very large T_{wd}), and a surface gravity range, log $g = 7.0 - 9.0$, in increments of 0.2 in log g.

When available, we adopted published E(B-V) values to de-redden the spectra. We took the published parameters for the six systems to use as initial estimates in our model fitting. The emission line regions were masked out in the model fitting. We carried out a range of disk and photosphere model fits to derive accretion rates, possibly estimate hot component masses, orbital inclinations, and to compare model-derived distances with distance estimates published in the literature. Our main focus is to determine whether an accretion disk is present, estimate the temperature of the white dwarf, estimate the accretion rate, and gain possible insights into the underlying mechanism of their outbursts. The underlying, accreting white dwarf could be hidden from us by the accretion disk (if a disk can form in a wind-accreting system) or by the nebula and red giant's wind. It is because of the multi-component flux contributions of these emitting sources that actual model fitting analyses of the hot components have been very few and daunting.

4. Model Fitting

For each spectrum, we determined the best-fitting single temperature white dwarf model and the best-fitting accretion disk-only model. Depending upon the success of these fits, we tested whether a combination of a disk plus a white dwarf significantly improved the fit. Using two chi square minimization routines, either IUEFIT for disks-alone and photospheres-alone or DISKFIT for combining disks and photospheres or two-temperature white dwarfs,

reduced χ^2_ν (χ^2 per degrees of freedom ν) values and a scale factor were computed for each model or combination of models. The scale factor, S, normalized to a kiloparsec and solar radius, can be related to the white dwarf radius R through: $F_{\lambda(obs)} = SH_{\lambda(model)}$, where $S = 4\pi R^2 d^{-2}$, and d is the distance to the source. For the white dwarf radii, we use the mass-radius relation from the evolutionary model grid of Wood (1995) for C-O cores. The best-fitting model or combination of models was chosen based not only upon the minimum χ^2_ν value achieved, but the goodness of fit of the continuum slope, the goodness of fit to the observed Lyman Alpha region (when not contaminated with emission) and consistency of the scale factor-derived distance with other distance estimates in Table 1.

5. System by System Discussion and Modeling Results

The fitting results are tabulated in Table 3 where for each system, by row, we list (1) the system name; (2) SWP spectrum ; (3) white dwarf mass (in solar masses); (4) the accretion rate in solar masses per year; (5) inclination in degrees; (6) model-derived distance (pc) (7) the reduced χ^2_ν value of the best fit; (8) the per centage of FUV flux due to accretion disk light and; (9) the percentage of FUV flux due to the white dwarf photosphere.

5.1. LT Del

LT Del has a yellow giant donor and has had only one recorded outburst. Skopal (2005) carried out an analysis of LT Del's single SWP spectrum along with a few optical flux points from the simultaneous optical spectroscopy by Munari & Buson (1992) who derived an accretion rate for LT Del of $\dot{M} = 3.0 \times 10^{-11} M_\odot$/yr. The steeply sloping continuum and high ionization emission line spectrum of LT Del suggests a high temperature for the hot component and a low amount of nebular material. Emission lines are strong with very high ionization degrees, and they include C IV (1550 Å), Si IV (1398 Å), N V (1240 Å), O IV] (1402 Å), He II (1640 Å), N IV (1485 Å), and O III (1662 Å). Additionally, there is an emission feature at 1498Å that is unidentified. The spectrum also lacks absorption lines that are usually found in hot stars, and it is also missing several lines that are common in symbiotics (Munari & Buson 1992).

With the IUE spectrum de-reddened with E(B-V) = 0.08, we attempted single temperature white dwarf model photosphere fits but this effort yielded model-derived distances that are far too close (due to their smaller stellar radii). Comparing the observed FUV continuum with white dwarf model atmospheres, the slope of the continuum agrees with a surface

temperature in the lower 20,000Ks (the exact value depending on Log g). For log g = 8.0 the best fit is for T=25,000K, d=138pc, $\chi^2 = 3.19$. For log g=9.0, the corresponding best-fit has T=27,000K, but the implied distance is only d=72pc, with $\chi^2 = 3.51$ while for a lower gravity, log g=7.4, T=25,000K, d = 200pc, and $\chi^2 = 3.00$ with $R_{wd} = 0.01974 R_\odot$. In Fig.1, we display this fit (log g = 7.4). Clearly, single temperature WD photosphere models cannot be correct because the implied distances are far too close. Moreover, the WD temperatures are unexpectedly cool for a WD in a wind accreting symbiotic system (Muerset et al. 1991) This conclusion points to an accretion disk as the most likely contributor to the FUV flux.

We found the best-fitting model ($\chi^2_\nu = 2.7$) to the spectrum of LT Del, to be an accretion disk with inclination i=60deg, an accretion rate to be $\dot{M} = 1.0 \times 10^{-8} M_\odot$/yr, a white dwarf mass of $1.0 M_\odot$ and a distance of 1450pc. This best-fitting accretion disk model is displayed in Fig.2. Skopal (2005) carried out an analysis of LT Del's single SWP spectrum along with a few optical flux points from the simultaneous optical spectroscopy by Munari & Buson (1992). He adopted a temperature for the hot component of 100,000K as its "characteristic temperature". However, it remained unclear whether a disk, white dwarf or both contribute to the FUV flux. The accretion rate that we have derived would certainly be high enough to heat the hot component to such a high temperature if the stellar component is truly that hot. In our disk model, the inner disk itself reaches a temperature of 93,000K. We note that Munari & Buson (1992) derived an accretion rate for LT Del of, $\dot{M} = 3.0 \times 10^{-11} M_\odot$/yr based upon black body fits to the FUV continuum slope and also found agreement with a hot component having a surface temperature of 200,000K. However, their analysis used an E (B-V) = 0.5!

Ellipsoidal light curves indicate a Roche Lobe-filling giant (Munari 2012), which suggests a much larger (a factor of a few or even more) distance then that indicated by our best fit. Indeed, for the systems with reported ellipsoidal variability, LT Del, BD-123873, and RW Hya, the distances in Table 1 are underestimated.

5.2. IV Vir (= BD -21 3873)

BD -21 3873 has a yellow giant donor and has never had a recorded outburst. The system has an orbital period of 281.6d and orbital parameters and masses from a combined radial velocity curve and light curve orbital solution by Smith et al. (1997). They derived a WD mass of $0.45 M_\odot$, semi-major axis a = 0.93 AU, and inclination i = 68 degrees.

The giant component is clearly metal-poor ([Fe/H]=-1.3), and s-process enhanced (Smith et al. 1997) due to pollution by mass transfer when the present white dwarf was a thermally puls-

ing asymptotic giant branch (TP-AGB) star. From the theoretical minimum core mass for such a TP-AGB star, the WD mass must be $> 0.55 M_{\odot}$ (Herwig 2005).

The ellipsoidal variability reported by Smith et al. (1997) is consistent with a Roche lobe filling giant. The distance set by the Roche lobe geometry of the red giant ($R_g = RL = 0.4au$) would be ∼4.4 kpc. However, if the giant is synchronized with the orbital period, then its rotational velocity, $Vsin(i)$ ∼6 km/s (Smith et al. 1997), thus implying a much smaller giant radius, Rg∼ $39 R_{\odot}$ (0.18 au), and a distance of no lower than 2 kpc. So, either the rotation in subsynchronous or the giant does not fill its tidal lobe (which is inconsistent with the ellipsoidal light curve).

The two IUE FUV spectra show continua steeply rising toward shorter wavelengths on which are superposed emission features due to N V (1240), Si IV (1400), an unidentified emission line at 1498Å, a strong C IV (1550) emission feature, He II (1640) and O III] 1667 emission. Outside of these emission lines, the continuum is essentially featureless. BD -21 3873 exhibits a rapidly variable bi-polar outflow, yet previous work by Skopal (2005); Munari et al. (2001) revealed no evidence of an accretion disk and negligible nebular contributions to their FUV continuum fluxes.

The value of E(B-V), 0.09, that we adopt is slightly lower than the total galactic reddening, E(B-V)=0.12, in the line of sight to BD-21 3873. An analysis by Skopal (2005) assumed a hot component temperature of $T_h = 100,000$K, while Schmid & Nussbaumer (1993) derived a hot component temperature of 75,000K using the modified Zanstra Method. We tried to obtain the best fits from modeling the FUV spectrum alone. However, the slope corresponds to only a 24-25,000K WD. The best-fitting WD photosphere model for IV Vir has $T_h = 24000$K, assuming log(g)=7.6 ($M_{wd} = 0.44 M_{\odot}$) and a $\chi^2 = 4.90$ and yields a distance of only 86 pc. Higher T_{wd} and lower T_{wd} have slopes that substantially disagree with the IUE spectrum. For example, $T_{wd} = 30,000$K corresponds to a distance of only 150pc, while $T_{wd} = 50,000$K yields 380pc. Next, we tried model accretion disk fitting to the IUE spectrum. We restricted the range of the inclination angle to be between 60 degrees and 75 degrees (see Table 1). The best-fitting model accretion disk corresponds $\dot{M} = 1.0 \times 10^{-8} M_{\odot}$/yr, $R_{wd} = 0.01513 R_{\odot}$, $M_{wd} = 0.55 M_{\odot}$, $Log(g) \sim 7.80$, with a $\chi^2 = 2.31$, and a model-derived distance d=517pc. Unfortunately the distance resulting from our best fit is too close and incompatible with both the lower limit distance of 2 kpc mentioned above and with the distance of ∼1kpc estimated by Pereira & Mello (1997). If we had allowed for a larger M_{wd} and a smaller inclination, then a larger distance would be obtained. We display the best-fitting accretion disk model to the IUE spectrum of IV Vir in Fig.3.

Our study marks the first time that an accretion rate was derived for BD-21 3873 by comparing the FUV spectra with models of optically thick steady state accretion disks.

Given that this object is a yellow symbiotic which has a giant donor of earlier spectral type and possibly weaker wind outflow, it is surprising that the accretion rate is as high as the disk model fitting indicates. On the other hand, it is not so high given that the giant likely fills it Roche lobe as suggested by the ellipsoidal light curve. While our fits with white dwarf photospheres also gave good agreement with the observations, we have ruled them out as the source of the FUV spectra because the implied distances are absurdly close and the temperatures far too cool to be the actual white dwarf temperatures since the contribution from cooler components (nebular continuum from the red giant wind photoionized by the Hot WD and the cool component Skopal (2005) tend to flatten the continuum slope in the FUV Skopal (2005).

5.3. StHa190

StHa190 has a confirmed jet outflow despite some studies suggesting that an accretion disk is absent (Skopal 1995). Munari et al. (2001) determined an orbital period of 171 days through a detailed search for variability and periodicities performed on Tycho and W95 data as well as the radial velocities. Smith et al. (2001) found a period of 37-39 days determined through periodicity searches of their own radial velocity data. Schmid & Nussbaumer (1993) classified it a D' symbiotic system. Munari et al. (2001) found the cool giant component to be a G7 III from the 3300-9100 Å spectra and MKK standards in MZ01. The E(B-V) value and distance were determined from comparing the MHZ photometry with the intrinsic colors from and absolute magnitudes from with E(B-V)=0.10 and d=575-780pc. Munari et al. (2001) asserted that a mass loss rate of $5 \times 10^{-8} M_{\odot}$/yr is needed to sustain the observed jets emanating from StHa190. Smith et al. (2001) determined the radius of the giant component to be $8.3 R_{\odot}$ and the effective temperature of the red giant to be 5025-5300K from spectral synthesis. The two IUE FUV spectra reveal a strong continuum rising steeply toward shorter wavelengths, strong C IV (1550) emission, strong O III (1667) emission, N III (1750) emission and a strong forbidden nebular line at C III [1909].

For StHa190, white dwarf photosphere fits alone reveal that the continuum slope corresponds to $T_{wd} = 22,000$K for a white dwarf mass $M_{wd} = 0.55 M_{\odot}$ ($R_{wd} = 0.01481 R_{\odot}$) but yields a distance of only 21 pc! Obviously, any WD photosphere can be ruled out as the source of the FUV flux for any reasonable white dwarf radius. The stellar radius would have to be larger by a factor of 30 to yield a distance of 600 pc, which is close to the distance range in Table 1. This finding led us to explore accretion disk models. Unfortunately, even the best fitting accretion disk models also yield normalization distances that are far too close.

We tried disk models with accretion rates of 10^{-8}, 3×10^{-9} and $10^{-9} M_{\odot}$/yr which yielded

distances, respectively of only 186 pc, 106 pc and 54 pc respectively. In Fig.4, we display the best fitting accretion disk model corresponding to an accretion rate of $10^{-8}M_{\odot}$/yr but with a scale factor derived distance of 186 pc. Unfortunately, in order to obtain a larger distance, comparable to Schild et al. (1996), the radius of the accreting white dwarf must be increased, thus decreasing the depth of the accretor's potential well and hence worsening the model disk fit.There are no models, disk or photosphere, that provide a reasonable fit. Therefore, it cannot be ruled out that our model accretion disks and model WD photospheres do not apply to the hot component in StHa190. One possibility might be that the hot component is not a white dwarf.

5.4. CQ Dra

CQ Dra was originally thought to be a cataclysmic variable in binary orbit with a red giant (Reimers et al. 1988), but was later re-classified as a symbiotic variable. However, as discussed below, the cataclysmic variable interpretation of the hot component may be correct after all. The IUE FUV spectra of CQ Dra are dominated by strong emission lines of C III (1175), weak N V (1240) emission, weak Si IV (1398) emission, very strong C IV (1550) emission, with He II (1640) being notably absent. CQ Dra has a reliable HIPPARCOS parallax $\pi = 5.25 \pm 0.48$mas (van Leeuwen 2007). The orbital period for CQ Dra was determined to be 1703 days by Eggleton et al. (1989) from radial velocity periodicity searches. This is the longest orbital period of the six systems discussed in this work. By matching an emission bump from the giant companion, Skopal et al. (1992) was able to determine the temperature of the red giant $T_{eff} = 3700$K, with d=178 pc, and $R_g = 95R_{\odot}$. By analyzing the ultraviolet continuum, they infer a white dwarf temperature $T_{eff} = 110{,}000$-$145{,}000$K, with a radius $R_{wd} = 0.007 - 0.006R_{\odot}$. The wind mass-loss rate was determined in Wheatley et al. (2003), following the prescription of and was found to be of the order of $\sim 10^{-8}M_{\odot}$/yr.

For the IUE spectrum of CQ Dra, we first tried single temperature white dwarf fits. For the WD only, we have $M_{wd} = 0.55M_{\odot}$, $R_{wd} = 0.0143R_{\odot}$, and a best-fit temperature of only $T_{wd} = 17,500$K a $\chi^2 = 16$ with a distance of only 28pc! This, however, was the best fit WD model to the IUE spectrum. In order to achieve a distance of ∼200 pc, the WD radius must be increased by a factor of 6. If the flux is really due to the WD, the contribution of the cooler components cannot be neglected be it the nebula (continuum and line emission from the photoionized red giant wind, or the secondary).

Since a strong photometric modulation is seen, the inclination should be fairly high. Therefore, we tried inclination angles of 40, 69 and 79 degrees which yielded, respectively,

accretion rates of 10^{-9}, 3×10^{-9} and $10^{-8} M_{\odot}$/yr. The χ^2 values for the three disk fits were not, in a statistical sense, significantly different from each other. The continuum slopes of the theoretical spectra are a slightly too steep. However, for smaller $\dot{M}$ or lower inclinations, the implied distances become considerably smaller ($<$ 135pc) than the parallax distance of 200 pc. All the disk models that correspond to a distance of about 200pc yield a χ^2 of almost 20. Some disk models with a lower $\dot{M}$ give a χ^2 of 13 but a distance of only 100pc.

In view of the parameters derived by Reimers et al. (1988) including the closer distance of 100 pc, we also generated model fits using the following values: E(B-V)=0.15, $d = 100 \pm$ 40pc and inclination $i = 50 \pm 15$ deg. The slope of the FUV continuum seen in the IUE spec'tra is consistent with a cool WD component. Our synthetic spectral Fits with a WD alone indicate $T \sim 13 - 15,000$K, but with a distance of only $\sim$ 20pc. Moreover, the Lyα profile is not consistent with a cold WD. A model with a WD temperature of 25,000K indicates a distance of only 60 pc and the quality of the model fit is bad. A higher temperature WD fit (45,000K) results in an even worse fit with a corresponding distance d = 200 pc. For the range of allowable inclinations (40 to 60 degrees) from Reimers et al. (1988), lower χ^2 values are reached for accretion disk models with accretion rates in the range of 10^{-9} to $3 \times 10^{-10} M_{\odot}$/yr and the distances resulting from these fits range from 173 pc down to 59 pc.

To our advantage, we found a usable FUSE spectrum of CQ Dra in the MAST archive that provided a new thrust to our investigation of CQ Dra. The FUSE spectrum shows broad and strong emission lines of C III (977 & 1175) and O VI with some additional emission lines but curiously the Lyman (series) profiles of H are not clearly seen, making it more difficult to identify a disk versus a WD. In Fig.5, we display the FUSE spectrum with the strongest, most reliably identified line features. The overall shape of the reddened spectrum is fairly flat and there is not much interstellar absorption seen in the spectrum. Some of the sharp emission lines are probably due to terrestrial airglow and some of the sharp weak absorptions may also be either terrestrial or interstellar. While identifing lines in the FUSE range via comparison with FUSE spectra of other systems with accreting white dwarfs, we found that the FUSE spectrum of the hot white dwarf in the dwarf nova RU Peg during quiescence, is very similar to the FUSE spectrum of CQ Dra as seen in Fig.6 There is also a similarity between the IUE spectra of the two systems as displayed in Fig. 7. These similarities suggest that the hot component in CQ Dra may indeed be a cataclysmic variable as Reimers et al. (1988) originally claimed thus identifying CQ Dra as a triple system.

In order to obtain a wider FUV wavelength coverage with which to carry out synthetic spectral fitting, we spliced the FUSE range to the IUE wavelength range. To do this we found three IUE SWP spectra of CQ Dra that a had a low flux level comparable to the

FUSE flux level in the spectral regions where the two spectra overlap (e.g. the region of C III 1175 and adjacent continuum). The slope of the best fit theoretical WD spectrum to the combination of the FUSE and IUE showed good agreement with the combined spectra when $T_{wd} = 30,000$K to 40,000K. For 50,000K and higher, discrepancies arise at the longer wavelengths but the overall fit is reasonably good as shown in Fig.8. Some parts of the FUSE spectrum show better agreement with a WD at $T_{wd} = 30,000$K while at the shortest wavelengths in the FUSE spectrum, there is better agreement with a WD surface temperature higher than 50,000K. However, the rise seen at the shortest wavelengths could be due to overlapping broad emission lines seen merged together at the FUSE resolution. When we fit the FUSE spectrum with an accretion disk model, then for a distance of 100 pc (the Reimers et al. (1988) distance), we obtain $\dot{M} = 10^{-9.5} M_{\odot}$/yr for disk inclinations of 40 to 60 degrees. If $\dot{M} = 10^{-9} M_{\odot}$/yr, the distance is of order 250 pc and the fit is poor. The only decent accretion disk fit to the FUSE spectrum of CQ Dra requires $\dot{M} = 1 \times 10^{-8} M_{\odot}$/yr and corresponds to a distance of ∼950 pc. Hence we discard the possibility that the FUSE FUV flux is due to a disk.

5.5. V443 Her

The IUE FUV Spectra of V443 Her are dominated by a clearly defined continuum sloping upward toward the shorter wavelengths with emission lines of Si IV, C IV , He II + O I (1640), C III [1909] and with N V (1240) notably absent. V443 Her has never had a recorded outburst. Its multi-wavelength data was analyzed most recently by Skopal (2005) who ruled out an accretion disk but fit the FUV data with a hot black body SED at $T_{eff} = 10^5$K. Dobrzycka et al. (1993) derived the orbital period, an E(B-V) value of 0.15, an estimated distance of ∼ 2 kpc, and an inclination of approximately 30 degrees. An upper limit to the distance is set by the Roche lobe geometry, and is of ∼3-4 kpc whereas a lower limit, ∼2 kpc, can be get assuming a normal M5 III giant (see Dobrzycka et al. (1993) for details). The total galactic extinction, E(B-V) = 0.13-0.15. Other estimates of E(B-V) range from 0.15 (Dobrzycka et al. 1993) to 0.2 (Muerset et al. 1991). The inclination should be low ($i <$ 60degrees) as indicated by the relatively low amplitude of photometric and spectrophotometric variability (Dobrzycka et al. 1993).

For the cool component, Dobrzycka et al. (1993) found a spectral type M5, a mass of $\sim 2.5 M_{\odot}$, a radius of $\sim 110 R_{\odot}$ and an effective temperature of 3300K. For the hot component, Dobrzycka et al. (1993) estimated a mass of $\sim 0.4 M_{\odot}$, a radius of $\sim 0.15 R_{\odot}$, and an effective temperature of 80,000K.

V443 Her has never had a recorded outburst. Its multi-wavelength data was analyzed

most recently by Skopal (2005) who dismissed an accretion disk but fit the FUV data with a hot black body SED at $T_{eff} = 10^5$ K. Unfortunately, the reddening is uncertain which complicates the continuum slope fitting. Dobrzycka et al. (1993) derived a distance of 2 kpc and a low mass WD ($0.5M_{\odot}$) for the system. and most other authors adopt these same parameters.

First, we carried out our analysis for two values of the reddening found in the literature, E(B-V) = 0.15 and E(B-V) = 0.5. For the case of E(B-V) = 0.15, our fitting yields a reasonable best-fit model $M_{wd} = 0.6M_{\odot}$, an orbital inclination of 18 degrees with an accretion rate of $10^{-8}M_{\odot}$/yr, implying a distance of 595 pc. This best-fitting accretion disk model is shown in Fig.9. Lower accretion rates and higher inclination angles improve the spectral fits but the distance to V443 Her becomes even closer.

If E(B-V) = 0.5, then the best fitting model is a hot white dwarf photosphere with $T_{eff} = 70,000$K, log g = 7 but an implied distance of only 328 pc. The best-fit accretion disk to the FUV data has i = 80 degrees and an accretion rate of $10^{-9}M_{\odot}$/yr but the distance also becomes unacceptably close. High gravity photosphere models have $T_{eff} \sim 30,000$ K which is close to Skopal's black body temperature for the hot component but again the implied distance is too close. The excitation level of the nebular emission, requires a much higher temperature ($> 75,000$K) than the value estimated by Skopal (2005) and also by our model fits. It is possible that the hot component has a disk-like structure flared at the disk edge, thus screening the radiation from the hot inner disk and hot white dwarfSkopal (2005). But this would only be possible if the inclination is such that the disk rim is being viewed nearly edge-on. If, on the other hand, the inclination is lower, then there is an inconsistency between the temperature required by the nebular emission and the temperature of the hot component if one assumes that the emission lines are due entirely to photoionization by the hot component.

Allowing the inclination for V443 Her to be larger only decreases the distance given by the fit. There is no model that is even close to 1kpc, so that the accepted 2 kpc distance is far from being reached. We checked the entire WD mass range (from 0.35 to $1.21M_{\odot}$) and that gave shorter distances as well. In addition, even the models that are close to 800 pc have a slope that is too steep slope compared to the observed spectrum. Adding a WD to the disk fits does not help either as the WD contributes only a small fraction of the flux with the "temperature" of the observed slope being only 20,000K.

In summary, our analysis of V443 Her suffers from a fundamental failure: Regardless of the parameters we use, the resulting model-derived distances are below 500pc, instead of 2000pc. Once again, we ascribe this failure to achieve the right distance to the flattening of the FUV continuum slope due to the contribution of cooler flux components. The flattening

of the observed FUV slope makes the hot component mimic a much cooler photosphere or disk.

5.6. RW Hya

RW Hydrae (=HD 117970) is an example of an S-type symbiotic system, with the primary classified as a M2 III red giant. The orbital period is 370.2 days (Merrill 1950; Kenyon & Mikolajewska 1995), and the inclination is quite high, just sufficient for the system to undergo eclipses. Estimates of the white dwarf mass range from 0.3 to 0.6 $M_{\odot}$, while that of the red giant is between 0.5 and $2M_{\odot}$ (Schild et al. 1996; Kenyon & Mikolajewska 1995). For the distance of the system, we adopt the value based upon corotation of the red giant determined by Schild et al. (1996) of 670 +/- 100 pc. At this distance the total luminosity of the system is $\sim 700L_{\odot}$. RW Hya exhibits ellipsoidal variability (Rutkowski et al. 2007). The distance set by the Roche lobe geometry, which requires a filling factor above 0.9, is then 1.7 kpc (Mikolajewska et al. 2014).

An important feature of RW Hya for white dwarf accretion studies is that there is little or no evidence for an accretion disk. Its multi-wavelength data was analyzed most recently by Skopal (2005) who dismissed an accretion disk but fit the FUV data with a hot black body SED at $T_{eff} = 10^5$ K.

For our analysis of the IUE spectrum, we used the system parameters given in Table 1, i.e. $i > 70$ degrees, $M_{wd} = 0.5M_{\odot}$, E(B-V) = 0.10 and d = 670 - 820 pc. We considered three possibilities in our modeling; an accretion disk, a single temperature white dwarf alone and a combination of a white dwarf plus an accretion disk.

For the IUE spectra, the distances implied by the single WD fits are far closer than 670 pc. The best WD model to the IUE continuum slope is only 26,000K and an implied distance of 30 pc. This WD fit is displayed in Fig.10. For a white dwarf mass of $0.4M_{\odot}$, even a surface temperature of 60000K corresponds to a shorter distance (160 pc) than 670 pc. Therefore, to match the large observed flux after de-reddening ($\sim 4 \times 10^{-12}$) with a white dwarf alone, the system has to be very close. Therefore, if the distance is 670 pc, then the WD would contribute only a small fraction of the FUV flux.

The best accretion disk fits for a low mass white dwarf, even with an accretion rate of $10^{-8}M_{\odot}$/yr, gave a distance of only 90 pc. The best-fitting accretion disk model to the IUE spectrum is shown in Fig.11. Adding a hot WD with T = 60,000K to the accretion disk yields a normalization distance still below 200 pc.

To our surprise, we found a good FUSE spectrum of RW Hya in the MAST archive which had never been analyzed and allowed us to sample the FUV flux down to the Lyman Limit. The FUSE spectrum of RW Hya indicates that the hot component has a very high temperature, quite possibly an accretion disk since there there is no sign of the Lyman series from the WD. It is possible that the photospheric light of the WD is undetectable when the accretion rate is large and the disk hot and luminous.

We carried out single temperature hot WD model fits for a low mass WD to the FUSE spectrum of RW Hya and found a best fit corresponding to a white dwarf with a surface temperature of 160,000K which yielded a scale factor-derived distance of 811 pc confirming that a low mass white dwarf with its larger radius, and a very high temperature gives a distance to RW Hya which is within the error bars of the original Schild et al. distance. This best-fitting NLTE WD model atmosphere fit to the FUSE spectrum is shown in Fig.12

We found a HST GHRS spectrum of RW Hya in the MAST archive that has a similar flux level to the FUSE spectrum, though its flux level is modestly higher and needs to be scaled down to about 70%. In this way, we extended the wavelength coverage from 1700A down to the Lyman Limit.After dereddening the spectra with E(B-V)=0.10, we found the same solution for the combined spectrum (FUSE+GHRS) that we found for FUSE alone, namely a very hot and very small mass WD can provide the flux needed. The fit reveals an accreting white dwarf with $T_{wd} = 160,000$K, $M_{wd} = 0.4$, and a scale factor distance of the order of 800pc. The best-fitting single temperature white dwarf photosphere model to the FUSE + HST spectrum is displayed in Fig.13. An ac creting white dwarf at this very high temperature almost certainly sustains thermonuclear burning of the accreted hydrogen and should be a supersoft X-ray source if obscuration of the emitted X-rays by the red giant wind and dust is ineffective.

6. Discussion

The six S-Type symbiotic binaries that we chose for this study were selected for their relative "simplicity" compared with symbiotics containing Mira variables with their high wind mass loss rates, more complex nebulae and associated dust. Our approach was to confront their FUV spectra with NLTE accretion disk models and hot white dwarf model photospheres, thereby testing hot component temperatures derived from modified Zanstra techniques, and achieving our key objectives; the accretion rate, accretion efficiency, and the photospheric temperatures of the hot components in S-type systems. These parameters are crucial for understanding the evolution of symbiotic stars, including whether accretion disks form and dominate the FUV light, and whether the relationship between the long term

time-averaged accretion rate and the surface temperature of the accretors via compressional heating (Sion 1995; Townsley & Bildsten 2003), also holds for symbiotic hot components.

Our approach has met with mixed success. It appears that, for most symbiotics, including the "simpler S-Types", their FUV spectra obtained with IUE, and HST (FOS,GHRS,STIS) of the hot components cannot be modeled in the same way that accreting white dwarfs and disks in cataclysmic variables in dwarf novae and nova-like variables can be modeled. The chief problem is that the observed FUV continuum slopes are less steep due to the inclusion of flux contributions principally from the nebular continuum produced by the photoionized red giant wind. Hence, the hot components mimic a much cooler white dwarf or cooler accretion disk thereby lowering the derived temperature or accretion rate, while the true continuum slope of the hot component alone is much steeperSkopal (2005). In addition, while 3D numerical simulations (Mohamed & Podsiadlowski 2007; de Val-Borro et al. 2014) the accretion disks in symbiotics resemble optically thick, steady state disks, the scale of the symbiotic systems is vastly different than cataclysmics, and the accretion flows are more complex. However, our investigation underscores the critical importance of extending the FUV coverage of accreting white dwarfs in interacting binary systems down to the Lyman Limit. This coverage was key to our understanding of the nature of the hot components in CQ Dra and RW Hydrae, the only two systems for which FUSE spectroscopic coverage was available.

Acknowledgements

This work is supported by NASA Grants NNX13AF11G and NNX13AF12G to Villanova University and by undergraduate summer research support by the NASA Delaware Space Grant Consortium. JM has been supported by by the Polish NCN grant no. DEC-2011/01/B/ST9/06145. PG is thankfull to William (Bill) P. Blair for his kind hospitality in the Henry Augustus Rowland Department of Physics & Astronomy at the Johns Hopkins University, Baltimore, MD.

REFERENCES

Arkhipova, V.P. et al. 1994, AstL, 20, 623

Diaz, M. P., Wade, R. A., Hubeny, I. 1996, RMxAC, 4, 95

Dobrzycka, D., Kenyon, S., & Mikolajewska, J. 1993, AJ, 106, 284

de Val-Borro, M.; Karovska, M.; Sasselov, D. D.; Stone, J. M. in "Highlights of Spanish Astrophysics VIII, Proceedings of the XI Scientific Meeting of the Spanish Astronomical Society held on September 8-12, 2014, in Teruel, Spain)

Eggleton, P.P., Bailyn, C. D., & Tout, C. A. 1989, ApJ, 345, 489

Fitzgerald, M.P.1970, A&A, 4, 234

Herwig, F. 2005, ARA&A, 43, 435

Hric et al.1991, Contrib. Astron. Obs. Skalnat Pleso, Vol.21,303

Hubeny, I. 1988, Comput. Phys. Commun., 52, 103

Hubeny, I., & Lanz, T. 1995, ApJ, 439, 875

Kenyon, S., & Mikolajewska, J. 1995, AJ, 110, 391

Livio, M.& Warner, B. 1984, Observatory, 104, 152

Merrill, P. 1950, ApJ, 111, 484

Mikolajewska J., Galan C., Hinkle K. H.,Gromadzki M., Schmidt, 2014, MNRAS, 440, 3016

Mohamed,S.,& Podsiadlowski,P. 2007, in Proceedings of the 15th European Workshop on White Dwarfs ASP Conference Series, Vol. 372, Eds.Ralf Napiwotzki and Matthew R. Burleigh. San Francisco: Astronomical Society of the Pacific, p.397

Munari, U., & Buson, L. M. 1992, A&A, 255, 158

Munari, U., et al. 2001, A&A, 369, L1

Munari, U., 2012, JAAVSO 40, 572

Muerset, U.& Nussbaumer, H. 1994, A&A, 282, 586

Muerset, U., Nussbaumer, H., Schmid, H. M., Vogel, M. 1991, A&A, 248, 458

Pereira, C. B,. & Mello, G. F. P. 1997, AJ, 114, 2128

Reimers, D., Griffith, R.F. & Brown, A. 1988, A&A, 193, 180

Rutkowski, A., Mikolajewska, M., Whitelock, P.A., 2007, Baltic Astr. 16, 49

Schild et al. 1996, A&A, 306, 477

Schild, H., Boyle, S. J., Schmid, H. M.1992, MNRAS, 258, 95

Schmid, H.M., Nussbaumer, H. 1993, A&A, 268, 159

Schmidt-Kaler, Th.1982, in Landolt-Bornstein Ser., vol. 2, Astron.Astrophys(Sringer-Verlag: Berlin)

Sion, E.1995, ApJ, 476, 875

Skopal, A. 1995, Ap&SS, 224, 557

Skopal, A. 2005, A&A, 440, 995

Skopal. A., et al. 1992, CoSka, 22, 131

Smith, V. V., Cunha, K., Jorissen, A., & Boffin, H. M. J. 1997, A&A, 324, 97

Smith, V.V., Pereira, C.B., & Cunha, K. 2001, ApJ, 556

Townsley, D., & Bildsten, L.2003, ApJ, 596, L227

Van Leeuwen, L. 2007, A&A, 474, 653

Wade, R. A., & Hubeny, I. 1998, ApJ, 509, 350

Wheatley, P.J., Mukai, K., & Martino, D. 2003, MNRAS, 346, 855

Wood, M. 1995, in *White Dwarfs, Proceedings of the 9th European Workshop on White Dwarfs* Held at Kiel, Germany, 29 August - 1 September 1994. *Lecture Notes in Physics*, Vol.443, edited by Detlev Koester & Klaus Werner. Springer-Verlag, Berlin, Heidelberg, New York, 1995, p.41

This preprint was prepared with the AAS LaTeX macros v5.2.

Table 1. Physical and Orbital Parameters

System Name	Distance <pc>	P_{orb} <days>	Inclination <deg>	M_{hot} M_{odot}	a <AU>	E(B-V)
LT Del	-	478.5[a]	-	-	1.1[b]	0.2[a]
BD-21	1000[f]	281.6[d]	68[d]	0.45[d]	-	0.09[d]
StH$_\alpha$190	575-780[a]	171[a]	60[b]	0.55[b]		0.10[a]
CQ Dra	178[c]	1703[d]	-	0.55[b]	0.10[c]	0.0
V443 Her	2000[k]	594[k]	$\sim$30[k]	$\sim$0.46[k]	0.115[l]	0.15[k]
RW Hya	670[m]	370.2[m]	$>$70[m]	0.5[m]	-	$<$ 0.1[m]

[a]Skopal (2005)

[b]Munari & Buson (1992)

[c]Arkhipova et al. (1994)

[d]Smith et al. (1997)

[f]Pereira & Mello (1997)

[k]Dobrzycka et al. (1993)

[l]Hric et al. (1991)

[m]Schild et al. (1992)

Table 2. IUE Observation Log

System	LT Del	BD-21 387	StHa190	CQ Dra	V443 Her	RW Hya
SWP#	39089	40933	47698	24265	09122	05939
Obs Date	6/14/1990	2/26/1991	5/20/1993	10/22/1984	5/25/1980	7/25/79
Exp. Time	7025	1800	1800	2880	1200	900
Continuum	63	55	74	126	131	223
Background	26	18	19	23	48	27

Table 3. Multi-Component Model Fitting Results

System	LT Del	BD-21 3873	StHa190	CQ Dra	V443 Her	RW Hya
Spectrum	swp39089	swp40933	SWP47698	SWP24265	SWP09122	SWP5939
Mass	1.00	0.55	0.55	0.55	0.60	0.40
$Log(\dot{M})$	-8	-8	-8	-	-8	-
Inclination (deg)	60	60	60	-	18	-
Distance (pc)	1450	517	186	438	595	800
χ^2_ν	2.87	2.31	8.9	2.15	9.86	-
% Flux_{disk}	100	100	100	-	100	-
% Flux_{star}	-	-	-	100	-	100



Fig. 1.— Flux vs. wavelength plot for the IUE spectrum SWP39089 of LT Del with a white dwarf model atmosphere with $Log(g) = 7.4$, $R_{wd} = 0.01974 R_{\odot}$, T = 25,000K, $\chi^2 = 3.00$ and a model-derived distance of only d = 200 pc



Fig. 2.— Flux vs. wavelength plot for the IUE spectrum SWP39089 of LT Del fitted with an accretion disk with inclination i = 60 deg, an accretion rate of $\dot{M} = 1.0 \times 10^{-8} M_{\odot}$/yr, a white dwarf mass of $1.0 M_{\odot}$ and a distance of 1450pc.



Fig. 3.— Flux vs. wavelength for IUE spectrum SWP40919 of IV Vir (=BD-21 3873). The best-fitting model accretion disk corresponds to $R_{wd} = 0.01513 R_{\odot}$, i = 60 degrees, $M_{wd} = 0.55 M_{\odot}$, Log(g)∼7.80, with a $\chi^2 = 2.31$, an accretion rate of $\dot{M} = 1 \times 10^{-8} M_{\odot}$/yrand a model-derived distance d = 517pc.



Fig. 4.— The best-fitting model fit to the de-reddened IUE spectrum SWP47698 of StHa190 with $M_{wd} = 0.55 M_{\odot}$, an inclination $i = 60°$, an accretion rate of $10^{-8} M_{\odot}$/yr but with a scale factor-derived distance of 186 pc.



Fig. 5.— The FUSE spectrum of CQ Dra with the strongest, most reliably identified line features. The overall shape of the reddened spectrum is fairly flat and there is not much interstellar absorption seen in the spectrum.



Fig. 6.— The FUSE spectrum of the hot white dwarf in the dwarf nova RU Peg during quiescence, shown in comparison with the FUSE spectrum of CQ Dra.



Fig. 7.— A comparison of the IUE spectrum of CQ Dra and the white dwarf in the dwarf nova RU Peg during quiescence.



Fig. 8.— The slope of the best fit theoretical WD spectrum to the combination of the FUSE and IUE spectra of CQ Dra shows good agreement with the combined spectra when $T_{wd} = 35,000$K to 40,000K. Here we show the $T = 40,000$K model with $\chi^2 = 2.15$, the distance is d=438pc for $M_{wd} = 0.55 M_{\odot}$.



Fig. 9.— A best-fit model to the IUE spectrum SWP09122 of V443 Her with $M_{wd} = 0.6 M_{\odot}$, an orbital inclination of 18 degrees with an accretion rate of $10^{-8} M_{\odot}$/yr, implying a distance of 595 pc.



Fig. 10.— The best WD model to the IUE continuum slope of RW Hya is only 26,000K and an implied distance of only 30 pc.



Fig. 11.— The best accretion disk fit for a low mass white dwarf, with an accretion rate of $10^{-8} M_{\odot}$/yr, gave a distance of only 90 pc.



Fig. 12.— A single temperature hot NLTE WD model fit for a low mass WD to the FUSE spectrum of RW Hya with a best fit corresponding to a white dwarf with a surface temperature of 160,000K, which yielded a scale factor-derived distance of 811 pc confirming that a low mass white dwarf with its larger radius, and a very high temperature gives a distance to RW Hya which is within the error bars of the original Schild et al. distance.



Fig. 13.— The de-reddened spectra of RW Hya with E(B-V)=0.10, for the combined spectrum (FUSE+ HST GHRS). a very hot and very small mass WD can provide the flux needed. The fit reveals an accreting white dwarf with $T_{wd} = 160,000$K, $M_{wd} = 0.4$, and a scale factor distance of the order of 800pc.